


Alisa Mast, Dave Clow, Jill Baron, Greg Wetherbee







Loch Vale Watershed







Loch Vale Watershed














Foundations

- 31 years of continuous monitoring
- · Paleolimnology
- Experiments in field and lab to test cause and effect
- Modeling ecosystem processes and "what-if" scenarios
- Spatial comparisons in Colorado and across western US





Annual runoff tracks precipitation







But first! When data are taken at face value...





1984-2012	Bear Lake SNOTEL	Loch Vale NADP	Loch Outlet
Bear Lake			
SNOTEL		0.56	0.6
Loch Vale			
NADP	0.56		0.29
Lake Irene			
SNOTEL	0.74	0.48	0.53
Copeland			
SNOTEL	0.71	0.48	0.52
University			
Camp SNOTEL	0.76	0.54	0.57
Niwot SNOTEL	0.66	0.52	0.57
Niwot C1	0.73	0.5	0.42
Niwot D1	0.62	0.59	0.33

R² of annual precipitation amount





Bear Lake SNOTEL had higher correlations with ANY site other than Loch Vale NADP





1984-2012	Bear Lake SNOTEL	Loch Vale NADP	Loch Outlet
Bear Lake SNOTEL		0.56	0.6
Loch Vale NADP	0.56		0.29
Lake Irene SNOTEL	0.74	0.48	0.53
Copeland SNOTEL	0.71	0.48	0.52
University Camp SNOTEL	0.76	0.54	0.57
Niwot SNOTEL	0.66	0.52	0.57
Niwot C1	0.73	0.5	0.42
Niwot D1	0.62	0.59	0.33

R² of annual precipitation amount





Even discharge at Loch Outlet compared poorly with Loch Vale NADP





1984-2012	Bear Lake SNOTEL	Loch Vale NADP	Loch Outlet
Bear Lake SNOTEL		0.56	0.6
Loch Vale NADP	0.56		0.29
Lake Irene SNOTEL	0.74	0.48	0.53
Copeland SNOTEL	0.71	0.48	0.52
University Camp SNOTEL	0.76	0.54	0.57
Niwot SNOTEL	0.66	0.52	0.57
Niwot C1	0.73	0.5	0.42
Niwot D1	0.62	0.59	0.33

R² of annual precipitation amount





We upgraded the Loch Vale precipitation gage from a Belfort to a Noah IV gage in 2007




which resulted in a step change in precipitation amount unmatched by nearby sites





NOAH IV captures more pptn than Belfort and SNOTEL



Vertical lines: range of annual pptn at 6 Front Range SNOTEL sites

📕 : Loch Vale Belfort gage


: Loch Vale NOAH IV gage





Four years of co-location of old and new gages allowed for a correction factor






%: increases in pptn at Noah IV compared to Belfort








[NO₃⁻] and [NH₄⁺] in Loch Vale NADP precipitation ↑until ~2004









[NO₃-] and [NH₄+] in Loch Vale NADP precipitation ↑until ~2004

Similar pattern from 6 other Front Range NADP sites











[NO₃-] and [NH₄+] in Loch Vale NADP precipitation ↑until ~2004

Similar pattern from 6 other Front Range NADP sites






Total N (wet+dry) deposition: same pattern of tand subsequent








Stream [NO₃-]
and yield:
same trends
as precipitation







Why is there such a direct response?








Why is there such a direct response?















83% variability at
Andrews Cr.
(73% at The Loch)
explained by
precipitation
[N]+ PDI lagged 1 yr








83% variability at
Andrews Cr.
(73% at The Loch)
explained by
precipitation
[N]+ PDI lagged 1 yr











83% variability at
Andrews Cr.
(73% at The Loch)
explained by
precipitation
[N]+ PDI lagged 1 yr





N accumulates in low runoff years, is flushed in normal/high years.


And N accumulates in biomass






What is causing \(\bar{\psi} \) N deposition?

Loch Vale



Front Range NADP sites

















Long-term monitoring essential for trend and pattern analysis

N deposition has been declining since ~2010

In high elevation Loch Vale stream N tracks deposition (with a lag due to drought)

Water quality is improving in real time in response to lower N emissions and deposition





